


# Topic Area TU: Tobacco Use

## Objective TU-6: Percent Smoke Cessation During First Trimester of Pregnancy\* - 2020 Target: 30.0

Percent Smoke Cessation During First Trimester of Pregnancy\* by Race and Hispanic Origin, Pennsylvania Residents, 2009-2013



Smoke Cessation During First Trimester of Pregnancy*, Percent and Number for Selected Demographics, Pennsylvania Residents, 2009-2013										
	PENNSYLVANIA									
	2009		2010		2011		2012		2013	
	%	No.	%	No.	%	No.	%	No.	%	No.
Smoke cessation during first trimester of pregnancy* mothers 18-49	25.6	7,733	26.2	7,597	26.3	7,378	26.2	7,054	26.7	6,879
White mothers 18-49	26.0	6,388	26.6	6,271	26.3	6,024	26.1	5,724	26.4	5,448
Black mothers 18-49	19.9	724	20.8	728	22.3	774	22.1	748	24.5	683
Hispanic mothers 18-49	28.8	534	32.9	598	33.1	577	36.1	597	34.9	586
Asian/Pacific Islander mothers 18-49	48.8	100	47.9	80	47.5	85	42.3	63	43.7	45

\*Mothers counted as quitting smoking were those who smoked during the three months prior to the pregnancy and did not smoke during the first, second, and third trimesters.

NOTE: Hispanics can be of any race.

The above statistics and additional data for Healthy People 2020 objectives can also be found at [www.statistics.health.pa.gov](http://www.statistics.health.pa.gov) (select Healthy People).

DSU = Data statistically unreliable (fewer than 10 events) NE = No events